

Scotland's Rural College

Reducing soil erosion in smallholder farming systems in east Africa through the introduction of different crop types

Muoni, Tarirai; Koomson, Eric; Öborn, Ingrid ; Marohn, Carsten; Watson, CA; Bergkvist, Göran; Barnes, AP; Cadisch, Georg; Duncan, Alan

Published in:
Experimental Agriculture

DOI:
[10.1017/S0014479719000280](https://doi.org/10.1017/S0014479719000280)

First published: 17/09/2019

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

Muoni, T., Koomson, E., Öborn, I., Marohn, C., Watson, CA., Bergkvist, G., Barnes, AP., Cadisch, G., & Duncan, A. (2019). Reducing soil erosion in smallholder farming systems in east Africa through the introduction of different crop types. *Experimental Agriculture*. <https://doi.org/10.1017/S0014479719000280>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Reducing soil erosion in smallholder farming systems in east Africa through the introduction of different crop types

Muoni Tarirai^{1,2*}, Koomson Eric³, Öborn Ingrid^{1,4}, Marohn Carsten³, Watson Christine A^{1,5}, Bergkvist Göran¹, Barnes Andrew⁶, Cadisch Georg³ and Duncan Alan^{2,7}

¹ Crop Production Ecology, Swedish University of Agricultural Sciences, P.O. Box 7043, SE-750 07 Uppsala Sweden,

² International Livestock Research Institute, 30709 Naivasha Rd, Nairobi, Kenya

³ Hans Ruthenberg Institute, University of Hohenheim, 70593 Stuttgart, Germany

⁴ World Agroforestry, P. O. Box 30677-00100, United Nations Avenue, Nairobi, Kenya

⁵ SRUC, Craibstone Estate, Aberdeen AB21 9YA, United Kingdom

⁶ SRUC, West Mains Road, Edinburgh, EH9 3JG, United Kingdom

⁷ Global Academy of Agriculture and Food Security, The Royal (Dick) School of Veterinary Studies and The Roslin Institute, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, UK

*corresponding author: +254733493834, +263774311136, tarirai.muoni@gmail.com

Abstract

On low-input smallholder farms of Kenyan upland landscapes, erosion of nutrient rich topsoil strongly affects crop yields. Where maize (*Zea mays*) is intercropped on erosion-prone slopes, intercropping can potentially reduce soil erosion. The objective of this research was to quantify the contribution of crops and crop mixtures of different growth habits to erosion control and their influence on above ground biomass and earthworm abundance as indicators of soil function in smallholder farming systems under a bimodal rainfall pattern in Western Kenya. The experiment involved five treatments, namely maize (*Zea mays*)/common bean (*Phaseolus vulgaris*) intercrop (maize intercrop), maize/common bean intercrop plus Calliandra (*Calliandra calothyrsus*) hedgerows and Calliandra mulch (Calliandra), sole Lablab (*Lablab purpureus*), sole Mucuna (*Mucuna pruriens*) and groundnut (*Arachis hypogaea*) intercropped with maize (during the short rains). The experiment was conducted over three consecutive cropping seasons and the cropping system had significant effects on soil loss, runoff, water infiltration, earthworm abundance and above ground biomass and crop grain yield. The Calliandra treatment had the lowest runoff (11.6 – 17.2 mm ha⁻¹) and soil erosion (31– 446 kg ha⁻¹, per season) in all the seasons, followed by the Mucuna treatment. Lablab was affected by disease and showed the highest soil erosion in the last two seasons. Infiltration was highest in Calliandra treatment, and earthworm abundance was higher under Mucuna and Calliandra treatments (229 and 165 earthworms m⁻², respectively) than under other crops. Our results suggest that including sole crops of herbaceous species such as Mucuna, or tree hedgerows with mixtures of maize and grain legumes has the potential to reduce

runoff and soil erosion in smallholder farming. Additionally, these species provide a suitable habitat for earthworms which stabilize soil structure and macropores and thus potentially increase infiltration, further reducing soil erosion.

Key words: Calliandra, infiltration, intercropping, legumes, Mucuna, runoff

1. Introduction

Land degradation is a major problem in sub-Saharan Africa (SSA) leading to low agricultural productivity and a need for increased inputs and ultimately leading to land abandonment and loss of land for food production (FAO, 2010). It has been estimated that approximately 30 % of land in SSA is degraded due to erosion, nutrient mining, overgrazing and deforestation (Bindraban et al. 2012). Apart from being a threat to food security and human nutrition, surface runoff and soil erosion are major environmental concerns (Fenta et al., 2017). Estimates of crop yield loss caused by erosion in SSA vary but may be substantial on severely eroded soils. For example, maize (*Zea mays*) yield reductions related to erosion have been estimated at up to 59 and 66 % in Tanzania and Kenya, respectively (Okoba and Sterk, 2010). Yield losses result not only from loss of soil organic matter and nutrients but also reductions in moisture availability and rooting depth. The rate of erosion from conventionally ploughed land has been estimated to be 1 to 2 orders of magnitude greater than both rates of erosion under native vegetation and rates of soil formation (Montgomery, 2007) suggesting an urgent need to look for alternative practices that reduce erosion rates. Rates of soil erosion depend on the interaction of several variables

(Verheijen et al., 2009), such as erosivity or energy of the eroding agent (wind and water), ground cover and management, slope characteristics and soil properties (erodibility). Control methods for water-induced erosion therefore aim to manipulate both ground cover and soil structure to reduce surface-water flow rates and increase infiltration rates. Methods include covering the soil as much as possible, minimum tillage, contour ploughing, use of grass strips planted on contours, inclusion of shrubs/trees (agroforestry) and use of cover crops (Salako et al., 2006).

In much of east Africa, effective erosion control practices have been poorly implemented (Okoba and Sterk, 2010). Two reasons for poor implementation of erosion control measures in Kenya are lack of evidence of their technical efficiency and social acceptability in different areas (Sigunga and Wandahwa, 2015). Another suggested reason for low uptake of erosion control measures is high initial costs relative to economically beneficial yield gains in the short term (Nyangena and Köhlin 2008). One method of erosion control, which does not require large investments, is increasing crop diversity potentially through utilizing species mixtures within a single growing season and in increasing diversity within crop rotations. Legume species (herbaceous, grain and tree legumes) are among candidates for increasing crop diversity due to their high nutritional value as food and fodder and their ability to biologically fix nitrogen. Herbaceous legumes are plants with non-woody stems above ground (Mongkhonsin et al., 2019) and in this paper we refer to them as crops mainly used as cover crops and/or fodder crops. Farmers in east Africa generally practice intercropping e.g. maize (*Zea mays*) and common bean (*Phaseolus vulgaris*), which increases soil cover and improves resource use efficiency compared to sole cropped maize (Himmelstein et al. 2017). Herbaceous and tree legumes may also be incorporated into cropping systems to reduce soil erosion while also providing other products and services. They have the

potential to contribute to soil erosion control because they have fast growth rates, high biomass production and some of them are drought tolerant (Kaspar et al., 2011). Rapid growth rates provide groundcover that helps reduce the impact of raindrops and direct sunlight hence improving soil and water conservation (Kaspar et al., 2011). High biomass productivity also increases soil organic matter content which improves soil structure as well as earthworm density, which contributes to water infiltration and holding capacity (Jordán et al., 2010). Additionally, legumes provide a series of other functions potentially useful to smallholder farmers including provision of food, livestock feed, income and soil fertility improvement through biological nitrogen fixation (BNF) and addition of organic matter. Tree legumes also have other purposes within smallholder farms including providing a source of fuel or construction material. Different legume types can be incorporated into smallholder farming systems through intercropping, in crop rotations, and planted on farm boundaries and as hedgerows. Livestock is important in east Africa and contributes 20–30 % of national gross domestic product, hence they play a significant a role in farming livelihoods (EAFF, 2012). The use of herbaceous legume species such as Mucuna (*Mucuna pruriens*) and Lablab (*Lablab purpureus*) and grain legumes such as cowpea (*Vigna unguiculata*) and groundnut (*Arachis hypogaea*) could thus play a useful role as providing fodder, as well as improving and protecting the soil.

There is little research that has focused on the ability of different legume types to contribute to the control of soil erosion in smallholder farming systems in east Africa. Thus, the main objective of this study was to assess the efficiency of integration of different crop types (represented by herbaceous, grain or woody legumes) in preventing surface runoff and soil erosion compared to maize-common bean intercropping that represents the typical cropping system in the region. Soil erosion measurements using runoff plots have been successfully

used in several recent papers and that method was applied in this study (e.g. Thierfelder and Wall, 2009). We also assessed the effect of different crop types and crop mixtures (with legumes) on water infiltration and earthworm abundance as indicators of soil function, and total above-ground biomass produced. The hypotheses tested were i) the introduction of herbaceous or woody species can reduce soil erosion compared to maize-common bean intercropping, and ii) herbaceous species and mulching enhance water infiltration and earthworm populations compared to annual grain legumes intercropped with maize.

2. Materials and methods

2.1 Site description

The experiment was conducted on-farm in Rongo district, Migori County (00°77' S, 34°60' E; 1474 meters above sea level), in western Kenya. The area is characterized by a sub-humid climate and receives rainfall in a bimodal pattern, with approximately 1000 mm per annum (Fig. 1). In general, the rainy seasons last from March to July (LR) and from September to December (SR). Annual average temperature is about 20 °C. Predominant soils are *Acrisols* according to FAO classification and the land is generally sloping. The soil at the experimental site is a sandy clay loam (Table 1) and the slope is around 20 %. Common crops in the region include maize (*Zea mays L.*), sugarcane (*Saccharum officinarum*), tea (*Camellia sinensis*), coffee (*Coffea arabica*), banana (*Musa L.*) groundnut and common bean (Rapsomanikis, 2015). The main livestock species are cattle, goats, sheep and chickens. Farmers utilize both cropping seasons and generally prefer to intercrop maize and common bean for food security and efficient land utilization, and the average farm size is approximately two ha (Rapsomanikis, 2015). Land

preparation and weed management are usually done with ox-drawn mouldboard plough or hand hoes (Rapsomanikis, 2015). The study farm had been cultivated for more than 10 years with mainly cassava, maize, common bean and groundnuts.

2.2 Experimental design

The experiment was established early in 2016 during the long rain (LR 2016) season and was continued during the subsequent 2016 short rain (SR 2016) and long rain (LR 2017) cropping seasons. It was arranged according to a randomized complete block design (RCBD) with five treatments and three replicates.

The treatments were as follows, maize/common bean intercrop (maize intercrop); groundnut, as sole crop during LR and intercropped with maize during SR (groundnut) (following farmer practice); Lablab (*Lablab purpureus*) as a sole crop (Lablab); Mucuna (*Mucuna pruriens*) as a sole crop (Mucuna) and maize/common bean intercrop plus *Calliandra calothyrsus* hedgerow and leaf mulch (Calliandra).

Soil samples were collected with a soil auger (internal diameters 7 cm) at two depths (0-20 cm and 20-40 cm) just before planting in April 2016. Ten soil samples from each depth were bulked to give one sample per block and sub-samples of 500 g were saved for further analysis. Soil pH was determined in 1:2.5 soil and 0.01 M CaCl₂ suspension using a Labor-pH-meter (WTW GmbH). Total N and C were analysed using dry combustion (Flash EA 1112 Elemental Analyser). Soil texture was determined using the pipette method, after having removed the organic matter using 35 % hydrogen peroxide. Available K was extracted using Calcium-

Acetate-Lactate solution and measured by Inductively Coupled Plasma Optical Emission Spectrometry (Agilent 5100 ICP-OES). Available P was extracted using Bray No 1 extractant and analysed on spectrophotometer (Bechman UV/Vis Spectrophotometer DU @-640). For bulk density determination, 110 cm³ undisturbed cylinder cores were taken, the soil oven dried at 105 °C, and the bulk density calculated using the volume of the core.

The crops were sown after the first effective rains in all seasons, except for the LR 2016 season when crops were established a bit later. Establishment was delayed by a long dry spell after the runoff plot structures were constructed (at the beginning of the rainy season). Land was prepared using an ox-drawn mouldboard plough to a depth of approximately 20 cm, at the onset of the experiment to remove *Cynodon nlemfuensis* Vanderyst (African Bermuda-grass) which was common on the site and to loosen the soil for improved seed establishment. In the following seasons (September 2016 SR and March 2017 LR), land preparation was carried out with hand hoes (tilling depth approximately 20 cm) 2 weeks after harvesting the preceding crops. From the SR 2016 cropping season, 50 % of the harvested leaf and stem biomass in all treatments was retained in the respective plots and was uniformly spread soon after sowing the following crop, following recommendations from Mupangwa and Thierfelder (2014) and allowing the remaining crop residues to be used for feeding livestock or other purposes. Each main plot measured 12 m × 6 m (72 m²) and consisted of a bounded runoff plot measuring 12 m × 4 m (48 m²) in the centre with a 1 m buffer zone on both sides of the aluminium sheets (Muoni et al., 2018). At the start of the experiment, aluminium sheet boundaries were buried 0.20 m into the soil around the runoff plots to prevent water flow from adjacent plots or outside of the experimental area. At the bottom of each runoff plot, a triangular cross-section was constructed with a 5-cm diameter iron

pipe outlet connected to two 100 L tanks to collect runoff and soil sediments. The first tank had 6 equidistant levelled splitter outlets. One splitter outlet was then connected to a second tank (splitter tank), to account for the overflowing water from the splitter on the first tank.

All crops and the Calliandra hedgerows were planted across the slope. Maize was sown at 0.75 m \times 0.30 m to achieve 44 444 plants ha⁻¹. Common bean was sown in-between the maize rows, spaced at 0.20m between plants giving 66 666 plants ha⁻¹. Mucuna and Lablab were sown at 0.50 m between rows and 0.20 m between plants (100 000 plants ha⁻¹). Sole groundnut was spaced 0.45 m \times 0.15 m to achieve 148 148 plants ha⁻¹, for intercropping they were sown between the maize rows at the same in-row spacing (74 074 plants ha⁻¹). Fertilizer applications in all crops followed the recommended application rates in the region. All maize plots received 100 kg ha⁻¹ of Diammonium Phosphate (DAP) (18 kg N: 46 kg P₂O₅ (20 kg P) ha⁻¹) fertilizer at sowing. Sole legumes received 45 kg ha⁻¹ of DAP (8 kg N: 21 kg P₂O₅ (9 kg P) ha⁻¹). Thus, intercrops received 145 kg ha⁻¹ DAP. Reseeding in places of poor germination ('gap filling') was done approximately 2 weeks after initial sowing. All maize crops were top dressed, at four and seven weeks after sowing in two equal splits, using calcium ammonium nitrate fertilizer (26 kg N ha⁻¹). Hand weeding was carried out at least twice in each cropping season and the first and second weeding was carried out at four and seven weeks after sowing in each season followed by a third weeding when deemed necessary. Aphids and leaf eating caterpillars were controlled using Karate (active ingredient (a.i) Lambda-cyhalothrin), and fungal diseases on legumes were controlled using Redomil (a.i. metalaxyl-M plus mancozeb) at recommended application rates.

The Calliandra hedgerows were established using small seedlings spaced at 4 m \times 0.50 m (19 August 2016) in the beginning of the SR 2016 cropping season (seedlings were not available LR

2016). Each plot had three hedgerows that were placed at the top, middle and bottom of the plot. The seedlings received supplementary watering for the first three weeks to prevent wilting. The seedlings grew naturally until the end of the LR 2017 when they were cut at 0.60 m height. Calliandra mulch was imported from the nearby farms in the LR2016 before the hedgerows were established, and also in the following seasons because the seedlings in the plots were too small to provide sufficient quantities of plant material. The mulch was applied at 5 Mg ha⁻¹ soon after sowing in all seasons.

2.3 Field measurements

2.3.1 Runoff and soil erosion

Runoff and soil erosion measurements were carried out after each rainfall event that resulted in accumulation of water in the tanks. The second tank accounted 1/6 of the overflow from first tank. The total overflow was then added to the volume of water in the first tank. The total volume from all the rain events was added and converted to give runoff in m³ ha⁻¹.

All soil sediments collected from each tank were weighed on each sampling occasion. Thoroughly mixed soil sediments subsamples weighing approximately 500 g were collected from both tanks for oven drying. In cases where the soil sediments collected were below 500 g the whole sample was used. Soil sediments subsamples were oven dried at 105 °C until a constant weight was reached. The quantity of soil sediments from the splitter tank was also multiplied by 6 as in the case of runoff. Soil erosion is reported on a dry weight basis in kg ha⁻¹.

224

225 2.3.2 *Water infiltration*

226 Infiltration measurements were carried out during the SR 2016 and LR 2017. Water infiltration
227 was measured on 14 November 2016 (67 days after planting) and 5 May 2017 (65 days after
228 planting) during the SR and LR respectively, using a single ring infiltrometer measuring 5.08 cm
229 in diameter and 12.70 cm depth. The ring was driven 5 cm into the soil in an area cleared of plant
230 material. The infiltration was measured by pouring 107 mL of water into the ring and recording
231 the time taken for the water to infiltrate the soil. The procedure was repeated at three random
232 positions in each plot.

233

234 2.3.3 *Earthworms*

235 Sampling of earthworms was carried out at three sampling points in each plot on two occasions,
236 in November 2016 (SR 2016) and May 2017 (LR 2017), i.e. approximately 60 days after sowing
237 when there was adequate soil moisture. The sampling was done in the buffer zone outside the
238 main runoff plot to avoid soil disturbance that may have affected soil loss results. A metal frame
239 measuring 0.25 m × 0.25 m was randomly placed in the plot and all crop residues in the area
240 were removed. The soil was extracted to a depth of 10 cm and hand sorted for earthworms. Both
241 earthworms and soil were returned to the sampling point after counting the earthworms.

242

243 2.3.4 *Above ground biomass and crop grain yield*

Biomass data were collected from eight central rows \times 3 m long, of each crop. The total fresh weight of biomass from the net plot was weighed and a subsample (500 g) was oven dried at 80 °C for 48 hours. The subsample dry weight and fresh weight ratio was used to determine biomass dry weight in kg ha⁻¹. All three Calliandra hedgerows were pruned during the LR 2017 season at 0.60 m from soil surface in each plot and the leaves and stems in each plot were weighed separately. The average weight of leaves and stems in the three rows were calculated to give fresh weights of each plot. Stems and leaves sub samples of approximately 200 g were collected at each weighing. The sub-samples were oven dried at 80 °C for 48 hours. In intercropping treatments, total biomass included all harvested crop material including Calliandra in the appropriate treatment.

The maize grain yield was harvested on 8 central rows, each measuring 3 m. The fresh weight of all cobs in the 8 rows was measured immediately after harvesting and a sub-sample of 10 randomly selected cobs was taken for air-drying. The fresh and air-dry weights of the sub-samples were measured and grain yield was calculated at the recommended 12 % moisture content. For common bean and groundnut intercropped with maize, the crops were harvested between 8 central maize rows, also measuring 3 m each. Total fresh weight of the pods was measured in the field and a pod sub-sample of approximately 500 g was collected for air-drying. The grain yield for legumes was calculated at 9 % moisture basis. Sole Mucuna and Lablab grain data was collected using the same procedure as for common bean and groundnuts.

2.4 Statistical analysis

All data collected was subjected to heterogeneity of variance and normality tests and then the analysis of variance (ANOVA) was carried out, using Statistix 9 statistical package for personal computers, to assess the treatment effects on soil loss, runoff, water infiltration, earthworm populations, and total above ground biomass and grain yield (both maize and legumes) produced. The means of the three sampling points for earthworms and infiltration per plot were used in statistical analysis. Mean separation was carried out using the least significance difference (LSD) test at $P \leq 0.05$ on all significant data.

3. Results

3.1 Runoff, soil erosion and water infiltration

Cropping season had a significant effect on runoff. Runoff was higher during the LR 2017 and SR 2016 seasons than in the LR 2016 season (Fig 2). Also, runoff differed significantly between crop treatments in all three cropping seasons (Fig 2). The treatment Calliandra showed the lowest runoff in all seasons. Mucuna was the second most efficient crop in reducing surface runoff across the seasons, while the effects of the other crops were inconsistent. During the LR 2016 cropping season, runoff from the groundnut treatment was as low as from the Calliandra treatment whereas in SR 2016 and LR 2017 runoff under groundnuts was as high as with the worst performing crops. In the LR 2016, Lablab and Mucuna treatments had similar and intermediate runoff, whereas during the SR 2016 and LR 2017 Lablab was diseased and did not differ significantly from the maize-common bean (control) and the groundnut treatments. Both LR seasons had an extended mid-season dry spell when compared to the SR season (Fig 1).

286

287 Soil erosion differed by cropping season with SR 2016 having more soil erosion than LR 2016
288 and LR 2017 (Fig. 3). Soil erosion was also affected by treatments during the three cropping
289 seasons investigated in this study (Fig. 3). Soil erosion was lowest on the Calliandra treatment in
290 all three seasons, and the Mucuna treatment was similarly low during the first and the last
291 seasons ($<500 \text{ kg ha}^{-1}$). During the LR 2016, Lablab, Mucuna and Calliandra caused similar and
292 lowest soil erosion, while the maize intercrop led to the highest erosion. Overall, there was much
293 greater soil erosion in all treatments during SR 2016 (up to 6000 kg ha^{-1}) as compared to LR
294 2016 and LR 2017, except for Calliandra where erosion remained low. In the SR 2016 season,
295 maize intercrop, groundnut and Lablab had the highest soil loss, Mucuna caused intermediate
296 and caused Calliandra the lowest. During the LR 2017 season, soil erosion was relatively low
297 overall ($500\text{-}1500 \text{ kg ha}^{-1}$) and similar to LR 2016. Soil erosion was highest under Lablab
298 followed by maize intercrop. Calliandra, Mucuna and groundnut showed the lowest soil loss.

299

300 Treatments had a significant effect on water infiltration during the SR 2016 only (Table 2). The
301 highest water infiltration was observed under Calliandra and Mucuna treatments, while the
302 lowest was observed in Lablab treatment. Calliandra caused 154 % higher infiltration rate than
303 Lablab and 107 % higher than the maize intercrop treatment.

304

305 *3.2 Earthworm population*

The treatments had a significant effect on earthworm populations during the LR 2017 season only (Fig. 4). Mucuna and Calliandra caused similar, large earthworm populations compared to other treatments. The earthworm populations tended to be larger in Calliandra than in the other treatments also in SR 2016 ($P = 0.28$). Treatments with low earthworm populations had high surface runoff (Fig. 5a) and soil loss (Fig. 5b) during the SR 2016 and LR 2017 cropping seasons.

3.3 Above ground biomass and crop grain yield

The total biomass produced was larger and more affected by treatment in the SR 2016 season than in the LR 2016 (Table 3). In the SR 2016 season, the highest biomass was observed in groundnut/maize intercrop, Mucuna and Calliandra treatments, while Lablab produced the smallest biomass. In the LR 2017 season, Calliandra produced the largest total biomass followed by maize intercrop.

Treatment had a significant effect on maize and legumes grain yields during the SR 2016, where Calliandra treatment was larger than maize intercrop and groundnut treatments. Mucuna treatment produced a higher grain yield than other legume treatments (Table 3). During the LR 2017 season, treatments had a significant effect on legume grain yields only and the same trend was observed where Mucuna had the largest yield (Table 3). LR 2017 season had higher crop yields than SR 2016 in all treatments except for Lablab.

4. Discussion

Soil erosion control is a critical issue in smallholder farming systems in SSA. The washing away of organic matter and nutrient rich topsoil, accompanied by low fertilizer use, results in low crop yields. The use of intercrops including different types of legumes has the potential to improve soil fertility through reduction of soil erosion and addition of nitrogen through biological nitrogen fixation.

4.1 Runoff, soil erosion, water infiltration and abundance of earthworms

Mucuna and Calliandra caused low soil erosion and runoff during the study period. Although the first season trial was established late, this did not affect the results since the same trend of less soil erosion was observed from Calliandra and Mucuna treatments as from other treatments throughout the study period. Soil erosion observed during LR seasons in this study was generally low when compared to other studies, e.g. 6.9 Mg ha⁻¹ recorded under no-till plus legume intercrop in Zimbabwe 2005/06 cropping season (Thierfelder and Wall, 2009), >2 Mg ha⁻¹ under different crop and tillage management in Western Kenya (Ampofo et al., 2002) and 52 Mg ha⁻¹ year⁻¹ on bare soils during 2010/11 cropping season in Southern Africa (Paterson et al., 2013). Since the general level of erosion was small compared to other experiments conducted under similar conditions, the relatively small differences found in the present experiment can be expected to be of greater importance in situations as in the work referred to. Thus, the experiment supported the hypothesis that herbaceous and woody species can reduce soil erosion compared to maize-common bean intercropping. Increased infiltration was observed under Mucuna and Calliandra treatments, which also supports the hypothesis that herbaceous legumes and tree legumes with mulching

enhance water infiltration and earthworm populations compared to intercropping of annual crops and sole grain legumes.

Soil erosion, runoff and infiltration were probably reduced by the high soil cover provided by the crop canopies and residues (Montgomery, 2007; Thierfelder and Wall, 2009). Retention of crop residues increased soil cover at the onset of the season before crop establishment, which reduced raindrop impact, soil erosion and runoff. Raindrop energy loosens and displaces soil particles and this energy is dissipated by ground cover (Kaspar et al., 2011). Living and non-living mulch also diminishes the velocity of runoff which reduces the dispersion of soil particles from their original source thus reducing erosion (Kaspar et al., 2011). Mulching also increases surface roughness and thus reduces runoff velocity and increases ponding, giving the opportunity for increased water infiltration (Jordán et al., 2010). The greater above ground biomass production in treatments with *Mucuna* and *Calliandra* also increases litter and organic material in the soil, which improves aggregation of soil particles. The addition of mulch favored earthworm population build up, and the *Calliandra* leaves dropping during the season increased litter which may have improved moisture retention and feed availability for earthworms (Buchholz et al., 2017). More soil cover protects soil from direct sunlight and thus evaporation of surface water. This improves the retention of soil moisture, which is important for earthworm survival (Ivask et al., 2006). Both mulching and earthworm buildup contribute to more water infiltration (Jordán et al., 2010). As the *Calliandra* hedgerows grew, they provided additional ground cover which helped reduce runoff and soil loss. This could be the effect of changes in microtopography of the hedgerows (Lin et al., 2009) and earlier development of leaves at the beginning of the season, when the existing shrub roots can make use of residual soil moisture.

Establishing and pruning legume hedgerows for high quantities of fodder (or mulch) is labor intensive. Although tree and herbaceous legumes are valuable feed sources for animals and have been shown to increase milk production (Paterson et al., 2013), farmers are less willing to incorporate them because they lack experience in growing these crops. There is a very low interest in herbaceous legumes since farmers prefer growing grain legumes to provide food security for their families (Muoni et al., 2019). One way to increase interest in the inclusion of herbaceous or tree legumes would be to stress its usefulness as fodder in addition to the soil fertility enhancing and soil conservation aspects, but adoption is more likely in cases where livestock play an important role in income generation.

The maize intercrop had higher runoff and soil erosion than the other treatments during the LR 2016 cropping season, probably due to the long dry spell. Due to late establishment in that season, both maize and common bean suffered during the long dry spell and the vegetative growth period and thus produced little groundcover that could prevent runoff and soil erosion. Maize and common bean were more affected by moisture stress than Mucuna, Lablab and groundnut. More soil erosion and runoff were observed when the sole crop was in poor health, as we observed with Lablab in the seasons with pest issues. In such situations, crop mixtures will generally perform better than sole crops, because it is less likely that two or more crops will fail simultaneously, and the additional crop helps maintain groundcover (Rapsomanikis, 2015). In addition, crops such as Mucuna, Lablab and groundnut are all relatively drought tolerant and can provide ground cover even under harsh conditions.

4.2 Above ground biomass and crop grain yield

Crop biomass is crucial in smallholder farms for livestock feed, soil improvement, fuel and construction. Incorporating different crop types e.g. legumes in smallholder farms helps provide soil cover and livestock feed of better quality that helps improve soil fertility and livestock productivity. Groundnut intercropped with maize produced the largest biomass in SR 2016 season while Calliandra (with maize and common beans) produced the largest biomass in LR 2017 season. Although the Calliandra treatment produced high biomass, farmers need to be aware of the potential yield penalty resulting from competition for water, nutrients, light and space between crops and hedgerows, which can be reduced by regular pruning of hedgerows (De Costa and Surenthran, 2005). Biomass during the LR 2016 was lower than during other seasons because of late planting, and a long dry spell during the vegetative growth stage. When crop residues remain in the field soil organic matter can increase (Jordán et al., 2010) and this can increase biological activity and soil aggregate formation. Thus, legume crops that produce high biomass enable dual benefits of livestock feed and soil improvement, which improve farm productivity.

The maize intercrops produced high grain yields that may help farmers remain food secure. The Calliandra treatment had the largest maize yield during SR 2016 season due to the additional organic matter that was added by leaf mulch from the previous season. However, the Calliandra treatment had lower maize yield in the final season than maize intercrop treatment which may have been due to increased competition with Calliandra hedgerows and subsequent yield loss (De Costa and Surenthran, 2005). Competitive crops such as Mucuna, may be rotated with maize, which can help control soil erosion, improve soil fertility, provide livestock feed and provide food for farmers (Bonsu and Asibuo, 2013).

416

417 **5.0 Conclusions**

418 Incorporating a mixture of crop types in cropping systems has the potential to reduce runoff and
419 soil loss, increase earthworm populations and infiltration in smallholder farms. Based on the
420 results from this study, we conclude the following; use of a mixture of crop types including
421 herbaceous and woody species in cropping systems increases soil cover, which reduces the
422 impact of raindrops, as well as runoff and soil erosion. High soil cover can be attained in various
423 ways including intercropping (for example establishing Calliandra hedgerows in maize/common
424 bean intercrop) or crop rotations with herbaceous legumes that produce large amounts of
425 biomass (for example Mucuna). By including a mixture of crop types water infiltration can be
426 improved through increased soil cover and soil organic matter. Mucuna and Calliandra
427 treatments produced both high soil cover and above ground biomass than treatments with only
428 grain crops, which resulted in high infiltration rates and numbers of earthworms. Large
429 earthworm populations contribute to increase water infiltration through soil aggregate formation
430 and increased porosity. Inclusion of legume crops with high biomass production allows farmers
431 to use some biomass for protein rich livestock feed in integrated crop-livestock systems while
432 improving soil fertility.

433

434 **Acknowledgments**

435 We wish to acknowledge the German Academic Exchange Scheme (DAAD), Scotland's Rural
436 College (SRUC) and SLU for funding this research. We are also grateful to KALRO Kisii Centre
437 in Kenya for logistical support throughout the research period. We would like to thank Johannes

Forkman for his advice on statistical analysis. Samuel Obuche, thank you for supporting the conduct of this work on your farm. Marcos Lana and Petra Gwaka, thank you for comments on the manuscript. We also thank the anonymous review for constructive comments.

References

- Ampofu, E.A., Muni, R.K., Bonsu, M., 2002. Estimation of soil losses within plots as affected by different agricultural land management. *Hydrol. Sci. J.* 47, 957–967.
<https://doi.org/10.1080/02626660209493003>
- Bindraban, P.S. et al. 2012. Assessing the impact of soil degradation on food production. *Terrestrial systems* 4(5): p.478–488.
- Bonsu, P.O., Asibuo, J.Y., 2013. Rotational Effects of Legumes On Maize Yield *Int. J. Sci. Technol. Res.* 2, 222-227
- Buchholz, J., Querner, P., Paredes, D., Bauer, T., Strauss, P., Guernion, M., Scimia, J., Cluzeau, D., Burel, F., Kratschmer, S., Winter, S., Potthoff, M., Zaller, J.G., 2017. Soil biota in vineyards are more influenced by plants and soil quality than by tillage intensity or the surrounding landscape. *Sci. Rep.* 7. <https://doi.org/10.1038/s41598-017-17601-w>
- De Costa, W.A.J.M., Surethran, P., 2005. Tree-crop interactions in hedgerow intercropping with different tree species and tea in Sri Lanka: 1. Production and resource competition. *Agrofor. Syst.* 63, 199–209. <https://doi.org/10.1007/s10457-005-1090-8>
- EAFF, 2012. East Africa Livestock Strategy. East Africa Farmers Federation, Nairobi.
- FAO, 2010. The Status of Conservation Agriculture in Southern Africa: Challenges and Opportunities for Expansion. REOSA Tech. Brief 03.
http://www.fao.org/ag/ca/doc/FAO_REOSA_Technical_Brief3.pdf
- Fenta, A.A., Hiroshi, Y., Katsuyuki, S., Nigussie, H., Takayuki, K., Dagnenet, S., Kindiye, E., Sewale, B.A., 2017. Spatial distribution and temporal trends of rainfall and erosivity in the Eastern Africa region. *Hydrol. Process.* 31, 4555–4567.
<https://doi.org/10.1002/hyp.11378>
- Himmelstein, J., Ares, A., Gallagher, D., Myers, J., 2017. A meta-analysis of intercropping in Africa: impacts on crop yield, farmer income, and integrated pest management effects. *Int. J. Agric. Sustain.* 15, 1–10. <https://doi.org/10.1080/14735903.2016.1242332>

- 468 Ivask, M., Kuu, A., Truu, M., Truu, J., 2006. The effect of soil type and soil moisture on
469 earthworm communities. *Agric. Sci* 17, 7–11.
- 470 Jordán, A., Zavala, L.M., Gil, J., 2010. Effects of mulching on soil physical properties and runoff
471 under semi-arid conditions in southern Spain. *CATENA* 81, 77–85.
472 <https://doi.org/10.1016/j.catena.2010.01.007>
- 473 Kaspar, T.C., Singer, J.W., Hatfield, J.L., Sauer, T.J., 2011. The Use of Cover Crops to Manage
474 Soil, in: ACSESS Publications. Soil Science Society of America.
475 <https://doi.org/10.2136/2011.soilmanagement.c21>
- 476 Lin, C., Tu, S., Huang, J., Chen, Y., 2009. The effect of plant hedgerows on the spatial
477 distribution of soil erosion and soil fertility on sloping farmland in the purple-soil area of
478 China. *Soil Tillage Res.* 105, 307–312. <https://doi.org/10.1016/j.still.2009.01.001>
- 479 Mongkhonsin B, Nakbanpote W, Meesungnoen O, Prasad MNV 2019. Chapter 4 - Adaptive and
480 Tolerance Mechanisms in Herbaceous Plants Exposed to Cadmium. In: Hasanuzzaman
481 M, Prasad MNV, Fujita M (eds) Cadmium Toxicity and Tolerance in Plants. Academic
482 Press, pp 73–109
- 483 Montgomery, D.R., 2007. Soil erosion and agricultural sustainability. *Proc. Natl. Acad. Sci.* 104,
484 13268–13272. <https://doi.org/10.1073/pnas.0611508104>
- 485 Muoni, T., Barnes, A., Öborn, I., Watson, C.A., Bergkvist, G., Shiluli, M., Duncan, A., 2019.
486 Farmer perceptions of legumes and their functions in smallholder farming systems in east
487 Africa. *Int. J. Agric. Sustain.* <https://doi.org/10.1080/14735903.2019.1609166>
- 488 Muoni, T., Koomson, E., Öborn, I., Marohn, C., Watson, C.A., Bergkvist, G., Barnes, A.,
489 Duncan, A., 2018. Effect of legume-crop mixtures on runoff and soil loss in Africa.
490 *Advances in Legume Science and Practice Conference*. *Asp. Appl. Biol.* 138
- 491 Mupangwa, W., Thierfelder, C., 2014. Intensification of conservation agriculture systems for
492 increased livestock feed and maize production in Zimbabwe. *Int. J. Agric. Sustain.* 12,
493 425–439. <https://doi.org/10.1080/14735903.2013.859836>
- 494 Nyangena, W., Köhlin, G., 2008. Estimating Returns to Soil and Water Conservation
495 Investments: An Application to Crop Yield in Kenya. Discussion Papers dp-08-32-efd,
496 Resources For the Future. www.efdinitiative.org/sites/default/files/efd-dp-08-32.pdf
497 (accessed 18/04/2018)
- 498 Okoba, B.O., Sterk, G., 2010. Catchment-level evaluation of farmers' estimates of soil erosion
499 and crop yield in the Central Highlands of Kenya. *Land Degrad. Dev.* 21, 388–400.
500 <https://doi.org/10.1002/ldr.1003>
- 501 Paterson, D.G., Smith, H.J., van Greunen, A., 2013. Evaluation of soil conservation measures on
502 a highly erodible soil in the Free State province, South Africa. *South Afr. J. Plant Soil* 30,
503 213–217. <https://doi.org/10.1080/02571862.2013.861029>

504 Rapsomanikis, G., 2015. The economic lives of smallholder farmers: An analysis based on
505 household data from nine countries. FAO. Rome, Italy.

506 Salako, F.K., Kirchhof, G., Tian, G., 2006. Management of a previously eroded tropical Alfisol
507 with herbaceous legumes: Soil loss and physical properties under mound tillage. Soil
508 Tillage Res. 89, 185–195. <https://doi.org/10.1016/j.still.2005.07.010>

509 Sigunga, D., Wandahwa, P., 2015. Land and soil resources and their management for sustainable
510 agricultural production in Kenya: Current position and future challenges. Egerton J. Sci.
511 Technol. 11.

512 Thierfelder, C., Wall, P.C., 2009. Effects of conservation agriculture techniques on infiltration
513 and soil water content in Zambia and Zimbabwe. Soil Tillage Res. 105, 217–227.
514 <https://doi.org/10.1016/j.still.2009.07.007>

515 Verheijen, F.G.A., Jones, R.J.A., Rickson, R.J., Smith, C.J., 2009. Tolerable versus actual soil
516 erosion rates in Europe. Earth-Sci. Rev. 94, 23–38.
517 <https://doi.org/10.1016/j.earscirev.2009.02.003>

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532





Figure 1. Cumulative rainfall for all cropping seasons. LR 2016 x-axis is in days after planting (25 May 2016), for SR 2016 (planted 8 September 2016) and LR 2017 (planted 1 March 2017) its days after the first day of the month when season starts.



7

8 Figure 2. Effect of treatments on runoff during the 2016 long rains (LR 2016) and short rains
 9 (SR 2016), and the LR 2017 in Rongo district, Migori County, in Western Kenya. Groundnut
 10 was intercropped with maize during the SR 2016 season. Means with different letters in the
 11 same cropping season are significantly different from each other. Error bars are standard error
 12 of mean. LSD means least significant differences.

13

14

15

16

17


18

19

20

21


22



23

24

25 Figure 3. Effect of treatments on soil loss during the LR 2016, SR 2016 and LR 2017
 26 cropping seasons. Groundnut was intercropped with maize during the SR 2016 season. Means
 27 with different letters in the same cropping season are significantly different from each other.
 28 Error bars are standard error of mean. LSD means least significant differences.



30

31 Figure 4. Effect of treatments on earthworm population during the 2016 short rains (SR 2016)
 32 and 2017 long rains (LR 2017) cropping seasons in Rongo. Groundnut was intercropped with
 33 maize during the SR 2016 season. Means with different letters in the same cropping season are
 34 significantly different from each other. Error bars are standard error of mean. LSD means least
 35 significant differences.

36

37

38


39

40

41

42

43



44

45 Figure 5. Effect of earthworms in different treatments on a) runoff and b) soil loss during the SR
 46 2016 and LR 2017 season. Significance of the correlations was not tested because the data points
 47 in the regression were not independent. x- earthworms m^{-2} .

Table 1. Soil properties at the experimental site in Rongo district, Migori County, Western Kenya

| Depth (cm) | pH | Org C (%) | Total N (%) | C/N ratio | BD (g cm ⁻³) | Avail P (mg kg ⁻¹) | Avail K (mg kg ⁻¹) | Sand (%) | Silt (%) | Clay (%) |
|------------|-----|-----------|-------------|-----------|--------------------------|--------------------------------|--------------------------------|----------|----------|----------|
| 0–20 | 4.8 | 1.0 | 0.1 | 9.9 | 1.3 | 0.9 | 61.0 | 63 | 12 | 25 |
| 20–40 | 4.9 | 0.9 | 0.1 | 10.0 | 1.4 | 0.1 | 77.0 | 56 | 13 | 31 |

pH (measured in 0.01M CaCl₂ extraction with soil to extraction solution ratio of 1:2.5); Org C = organic carbon, N = nitrogen, BD=bulk density, Available phosphorus (P) and potassium (K). Available P was determined by Bray 1 with Beckman coulter Du, UV – Du 640 spectrophotometers, USA. Available K was analyzed by Calcium–Acetate–Lactate–extraction method.

Table 2 Effect of treatments on infiltration in Rongo during the short rains in 2016 and long rains in 2017

| Treatments | Short rains 2016 | Long rains 2017 |
|--|--|--|
| | Infiltration (mL second ⁻¹) | Infiltration (mL second ⁻¹) |
| Maize/Common bean intercrop | 2.7 ^b | 1.9 |
| Groundnut* | 3.0 ^b | 2.0 |
| Lablab | 2.2 ^b | 1.3 |
| Mucuna | 5.0 ^a | 1.4 |
| Calliandra | 5.6 ^a | 2.6 |
| Least significant differences (LSD) | 1.55 | NS |
| P-Value | 0.003 | NS |

*Groundnut was intercropped with maize during the SR 2016 season. Means with different letters are significantly different from each other.

Table 3 Crop grain and above ground biomass yield (kg ha⁻¹) during the short rains 2016 and long rains 2017 in Rongo. Statistical significances are indicated with different letters (p<0.05) for maize^(A, B) and legume^(a, b) grain yield. Above ground biomass yield is the total biomass of the treatment.

| Treatments | Crops | SR 2016 | LR 2017 | LR 2016 | SR 2016 | LR 2017 |
|-----------------|-------------|------------------------------------|-------------------|---|--------------------|--------------------|
| | | Grain yield (kg ha ⁻¹) | | Above ground biomass yield (kg ha ⁻¹) | | |
| Maize intercrop | Common bean | 79 ^b | 229 ^b | 3192 | 4336 ^B | 8532 ^A |
| | Maize | 3071 ^B | 7260 | | | |
| Groundnut | Groundnut | 73 ^b | 418 ^b | 2411 | 6592 ^A | 3040 ^B |
| | Maize | 3361 ^B | - | | | |
| Lablab | Lablab | 209 ^b | 80 ^b | 2318 | 548 ^C | 176 ^C |
| Mucuna | Mucuna | 1626 ^a | 3433 ^a | 2963 | 6197 ^{AB} | 5097 ^B |
| Calliandra | Common bean | 108 ^b | 240 ^b | 3481 | 5257 ^{AB} | 10426 ^A |
| | Maize | 4339 ^A | 4697 | | | |

‘LR 2016 has no grain yield data due to poor crops/crops not reaching maturity. Groundnut was intercropped with maize during the SR 2016 season and grown as sole crop LR 2017 (following farmer practice).